

1. Record Nr.	EJ21001523
Autore	Filipe, Joel Fernando Soares
Titolo	Rumen fluid, a new diagnostic matrix in dairy cattle farms? [Articolo]
Editore	Department of Health, Animal Science and Food Safety (VESPA), 2016-06-06
ISSN	2283-3927
Altri autori (Persone)	Trevisi, Erminio Massara, Matteo Minuti, Andrea Bani, Paolo Amadori, Massimo Riva, Federica
Lingua di pubblicazione	Inglese
Formato	Articoli digitali
Livello bibliografico	Seriale
Note	Diritti: Copyright (c) 2016 International Journal of Health, Animal Science and Food Safety In relazione con: https://riviste.unimi.it/index.php/haf/article/view/7039/7053 Sorgente: International Journal of Health, Animal Science and Food Safety; V. 3 N. 1s (2016) Sorgente: International Journal of Health, Animal Science and Food Safety; Vol 3 No 1s (2016) Sorgente: 2283-3927
Sommario	Production diseases of dairy cows are considered man-made problems caused by the inability of cowsto achieve a sufficient feed energy intake (Mulligan, 2008).A correct management of production diseases demands early diagnostic and prognostic parameters, in order to improve the management system and reduce the prevalence of clinical cases (Ingvarsen,2003).A previous study of our group indicated that forestomachs walls express immune receptors andcytokines, and the rumen liquor contains leukocytes able to produce IFN- (Trevisi, 2014).Our working hypothesis implied

that ruminal fluids could be a source of diagnostic information for the identification of herds at risk for production diseases. We first demonstrated that the diet can influence the immune response in forestomachs. Diverse leukocyte populations at low concentrations and IFN- were revealed in some samples of rumen fluids, with a clear inhibition of the response observed in the animals fed the maize-supplemented diet, compared to a normal and a soy-supplemented diet. We better characterized the leukocytes subpopulations in the rumen liquor, isolating B cells, monocytes and T cells. Finally we performed a field survey in order to find correlation among the immune profile of the rumen liquor. Clinically healthy animals showed a farm specific immunologic pattern of the rumen liquor: low CD45 mRNA expression, low IFN-, few/absent B-cells. We can conclude that the epithelial cells of ruminant forestomachs can react to different stresses (metabolic, infectious, inflammatory) and the inflammatory response can be sustained by infiltrating leukocytes. Our data points into the idea that dairy farms could be ranked according to a risk score using the inflammatory markers in rumen fluids, in addition to the traditional analysis.

Localizzazioni e accesso

http://memoria.depositolegale.it/*/https://riviste.unimi.it/index.php/haf/article/view/7039
